

Rempel et al. 2012: Depth and Chinook HSI (Fraser River)

Downloaded on: 2026-04-09, From: <https://mjbayly.com/stressor-response/rempe-et-al-2012-depth-and-chinook-hsi-fraser-river>
Function Updated by stobias on Mon, 12/23/2024 - 16:28.

Species Information

Common Name: Chinook Salmon
Genus: *Oncorhynchus tshawytscha*

Stressor Details

Stressor Name: Depth
Units: m
Metric: Water depth
Scale: linear
Function Type: continuous

Life Stage & Context


Life Stages: Fry
Geography: Fraser River

Descriptions

Overview

A wide range of depths were suitable for 0+ stream/ocean-type Chinook Salmon captured in spring (Figure 21). The wide depth suitability may be reflective of both shallow water use by stream-type fry and a broader distribution of migrating ocean-type fish. Depth suitability for Fraser gravel reach fish is similar to the WUP curve for Chinook fry, except that suitability declines past 80 cm for Fraser fish (Figure 21).

Stressor Response Data



Stressor (X)	Mean System Capacity (%)	SD	low.limit	up.limit
0	0	0	0	0
0.25	100	0	100	100
0.8	100	0	100	100
0.85	80	0	80	80
1	55.08	0	55.08	55.08

Citations

Rempel, L. L., Healey, K., & Lewis, F. J. A. (2012). Lower Fraser River juvenile fish habitat suitability criteria. Ecosystem Management Branch, Fisheries and Oceans Canada.

References

Rempel 2012 - <https://www.ecofishresearch.com/wp-content/uploads/2016/09/346413.pdf>