

Naman 2026: HSI Coho Juvenile Velocity (6°C daily net energy intake)

Downloaded on: 2026-10-16, From: <https://mjbayly.com/stressor-response/naman-2026-hsi-coho-juvenile-velocity-6degc-daily-net-energy-intake>
Function Updated by mbayly on Thu, 10/15/2026 - 15:58.

Species Information

Common Name: Coho Salmon
Genus: *Oncorhynchus kisutch*

Stressor Details

Stressor Name: Velocity
Units: m/s
Metric: Water Velocity (focal velocity) at 6°C
Scale: linear
Function Type: continuous
Vital Rate/Process: Habitat Suitability Index

Life Stage & Context

Life Stages: Fry, Parr
Geography: British Columbia (composite BC invertebrate database)
Activity: rearing
Season: Summer rearing

Descriptions

Overview

Temperature-dependent velocity suitability for a 6.5-cm, 3-g juvenile Coho Salmon, derived from the daily net energy intake (NREI_d, J/d; black line) predicted by a drift-foraging bioenergetic model at 6°C. NREI_d = daily gross energy intake (capped by temperature-dependent maximum consumption, C_{max}) minus daily foraging and standard metabolic costs. Values here are scaled 0–100 relative to the maximum of this 6°C line only (peak ~1104 J/d at ~0.07 m/s), not the global maximum across temperatures used for the paper's Figure 3 HSCs. At 6°C gross intake is constrained by C_{max}, producing a plateau and a low optimal velocity. Points were taken at the plotted model vertices (0.01–0.33 m/s, every 0.02 m/s) from the figure's vector line work; a final 0 was added at 0.340 m/s by extrapolating the last segment to NREI = 0, since the paper assigns 0 suitability to negative energy intake.

Stressor-response relationship data sourced from: Figure 2 — daily net energy intake (black line) vs. velocity, 6°C panel (left) (p. 4).

See linked SR Curve for 12C:
<https://mjbayly.com/stressor-response/naman-2026-hsi-coho-juvenile-velocity-12degc-daily-net-energy-intake>

Function Derivation

Mechanistic drift-foraging (NREI) bioenergetic model expressed on a daily time step with temperature-dependent costs and C_{max} (BioenergeticHSC + Wisconsin bioenergetics).


Transferability of Function

Specific to a 6.5-cm Coho at the stated temperature and BC composite drift; conceptual worked example rather than a site-calibrated curve.

Source of Stressor Data

Model output (BioenergeticHSC) using composite BC invertebrate drift data, 14 h foraging time, 50% reduction in prey detection/capture.

Stressor Response Data



Raw Stressor Values	Scaled Response Values 0 to 100	SD	low.limit	up.limit
0.01	9	0	0	100
0.03	44	0	0	100
0.05	77	0	0	100
0.07	100	0	0	100
0.09	98	0	0	100
0.11	96	0	0	100
0.13	93	0	0	100
0.15	90	0	0	100
0.17	86	0	0	100
0.19	82	0	0	100
0.21	79	0	0	100
0.23	74	0	0	100
0.25	70	0	0	100
0.27	58	0	0	100
0.29	41	0	0	100
0.31	24	0	0	100
0.33	8	0	0	100
0.34	0	0	0	100

Citations

Naman, S. M. (2026). Developing temperature-dependent habitat suitability curves for instream flow modeling. North American Journal of Fisheries Management. Advance online publication. <https://doi.org/10.1093/najfmt/vqag077>