

Ptolemy (n.d.): HSI Coho Juvenile Depth

Downloaded on: 2026-10-16, From: <https://mjbayly.com/stressor-response/ptolemy-nd-hsi-coho-juvenile-depth>
Function Updated by mbayly on Thu, 10/15/2026 - 16:05.

Species Information

Common Name: Coho Salmon
Genus: *Oncorhynchus kisutch*

Stressor Details

Stressor Name: Depth
Units: m
Metric: Water Depth (total depth)
Scale: linear
Function Type: continuous
Vital Rate/Process: Habitat Suitability Index

Life Stage & Context

Life Stages: Fry, Parr
Geography: British Columbia (composit)
Activity: rearing
Season: Summer (rearing)

Descriptions

Overview

British Columbia juvenile coho depth habitat suitability curve (R. Ptolemy, unpublished) used by Rosenfeld & Ptolemy (2012). Suitability rises steeply from 0 at 0 m to ~0.67 at 0.11 m and ~0.84 at 0.16 m, reaches 1.0 at ~0.25 m, and stays at 1.0 in deeper water (no deep-water decline). Values were extracted from the vector line work of the published figure (solid polyline vertices), so they approximate the original curves to within ~0.01 m and ~1 suitability unit. Note the figure's axis labels read $\text{cm} \cdot \text{s}^{-1}$ and cm, but the scales (0–1.5) are clearly m/s and m.

Stressor-response relationship data sourced from: Rosenfeld & Ptolemy (2012) Fig. 2f - Depth habitat suitability curve for coho salmon, Ptolemy (solid line) (p. 1923).

This curve seems to be used very frequently across instream flow studies, despite being unpublished. Users are encouraged to source their own data. There is no data associated with this relationship (context, fish age/stage/size, etc.).

Function Derivation

Empirical: fish density assessments coupled with stream transects (R. Ptolemy, unpublished data).

Transferability of Function


Developed from BC streams across a range of sizes; underlying data unpublished.

Source of Stressor Data

R. Ptolemy (BC Ministry of Environment), unpublished juvenile coho density and transect data from streams throughout British Columbia.

Stressor Magnitude Data

Stressor Response Data



Raw Stressor Values	Scaled Response Values 0 to 100	SD	low.limit	up.limit
0	0	0	0	100
0.113	67	0	0	100
0.16	84	0	0	100
0.252	100	0	0	100
1.5	100	0	0	100

Citations

Rosenfeld, J. S., & Ptolemy, R. (2012). Modelling available habitat versus available energy flux: Do PHABSIM applications that neglect prey abundance underestimate optimal flows for juvenile salmonids? *Canadian Journal of Fisheries and Aquatic Sciences*, 69(12), 1920–1934.

References

Rosenfeld & Ptolemy, 2012 - <https://doi.org/10.1139/f2012-115>