

Fine Substrate Particles and Kokanee Salmon Embryo

Downloaded on: 2026-04-08, From: <https://mjbayly.com/stressor-response/fine-substrate-particles-and-kokanee-salmon-embryo>
Function Updated by stobias on Sat, 11/16/2024 - 21:39.

Species Information

Common Name: Kokanee Salmon
Genus: *Oncorhynchus nerka*

Stressor Details

Stressor Name: Sedimentation
Units: % smaller than 6.35mm
Metric: Substrate particles
Scale: linear
Function Type: continuous

Life Stage & Context

Life Stages: Egg
Geography: Idaho

Descriptions

Overview


[Taken from figure caption] Relation between embryo survival and percentage of substrate particles smaller than 6.35mm for chinook salmon.

[Taken from paper] Redds that remain intact during incubation may become less suitable for embryos if inorganic fine sediments and organic materials are deposited in the interstitial spaces between the larger particles. The fine particles impeded the movement of water and alevins in the redd, and the organic material consumes oxygen during decomposition; if the oxygen is consumed faster than the reduced intragravel water flow can replace it, the embryos or alevins will asphyxiate.

Function Derivation

field study

Stressor Response Data



| Stressor (X) | Mean System Capacity (%) | SD | low.limit | up.limit |
|--------------|--------------------------|------|-----------|----------|
| 0.00 | 97.69 | 0.00 | 97.69 | 97.69 |

| | | | | |
|-------|-------|------|-------|-------|
| 5.99 | 95.05 | 0.00 | 95.05 | 95.05 |
| 9.63 | 93.07 | 0.00 | 93.07 | 93.07 |
| 12.89 | 90.10 | 0.00 | 90.10 | 90.10 |
| 17.44 | 86.14 | 0.00 | 86.14 | 86.14 |
| 20.69 | 81.85 | 0.00 | 81.85 | 81.85 |
| 24.34 | 75.58 | 0.00 | 75.58 | 75.58 |
| 26.68 | 70.63 | 0.00 | 70.63 | 70.63 |
| 30.59 | 61.06 | 0.00 | 61.06 | 61.06 |
| 34.88 | 48.84 | 0.00 | 48.84 | 48.84 |
| 37.74 | 41.25 | 0.00 | 41.25 | 41.25 |
| 40.48 | 33.33 | 0.00 | 33.33 | 33.33 |
| 43.08 | 27.06 | 0.00 | 27.06 | 27.06 |
| 45.68 | 21.12 | 0.00 | 21.12 | 21.12 |
| 48.81 | 15.18 | 0.00 | 15.18 | 15.18 |
| 52.45 | 10.23 | 0.00 | 10.23 | 10.23 |
| 56.75 | 4.95 | 0.00 | 4.95 | 4.95 |
| 59.87 | 3.30 | 0.00 | 3.30 | 3.30 |

Citations

Bjornn, T. C., & Reiser, D. W. (1991). Habitat requirements of salmonids in streams. American Fisheries Society Special Publication, 19(837), 99.

References

https://www.for.gov.bc.ca/hfd/LIBRARY/FFIP/Bjornn_TC1991.pdf -
https://www.for.gov.bc.ca/hfd/LIBRARY/FFIP/Bjornn_TC1991.pdf